

Research progress of oxidative stress in sepsis-associated liver injury

Hui Su^{1,2}, Tianying Xu³, Renke Sun^{1,2}, Yu Xiang^{1,2}, Yangmengna Gao^{1,2}, Kecheng Zai^{1,2}, Shangping Fang^{1,2}

¹School of Anesthesiology, ²Anesthesia Laboratory and Training Center, Wannan Medical College, Wuhu 241002, China. ³School of Anesthesiology, Department of Anesthetic Pharmacology, Second Military Medical University/Naval Medical University, Shanghai 200433, China.

Corresponding author: Shangping Fang.

Acknowledgements: This work was supported by Key Project Research Fund of Wannan Medical College (WK2022Z10) and Research Grant for College Students of Wannan Medical College (WK2022XS26).

Declaration of conflict of interest: None.

Received September 17, 2023; Accepted November 28, 2023; Published March 31, 2024

Highlights

- The approaches of antioxidant therapy to alleviate sepsis-associated liver injury are summarized from the perspective of oxidative stress in different cells of the liver.
- Reactive oxygen species, one of the main substances that induce oxidative stress, affects the molecular mechanism of the relevant signaling pathways.
- Antioxidant therapy is helpful for the recovery of various liver cells in sepsis-associated liver injury and is expected to advance basic and clinical research.

Abstract

Sepsis is a systemic inflammatory response that caused by infection or trauma, often resulting in multiple organ dysfunction. Its mortality rate is relatively high, ranging between 54% and 68%, and ineffective treatment and poor prognosis pose significant challenges to healthcare in recent years. One of the main pathogenesises of sepsis-induced liver injury is oxidative stress (OS), which refers to a state where the antioxidant system cannot balance oxidative products, leading to the accumulation of excessive oxidative products in the body. When sepsis occurs, the amount of reactive oxygen species produced by the body increases far beyond the levels that can be scavenged by the antioxidant system, thus damaging liver cells and aggravating liver damage. This article introduces the oxidative/antioxidant system, oxidative stress-related pathways, and the molecular mechanism of OS in various types of hepatocytes, with emphasis on the antioxidant treatment on different hepatocytes, in order to understand the mechanism of OS involved in the development and progression of sepsis-associated liver injury. As the research deepens, improving liver function through the treatment of different cells and facilitating related clinical research are expected to provide a new target pathway for the treatment of sepsis-associated liver injury.

Keywords: Sepsis-associated liver injury, liver injury, oxidative stress, reactive oxygen species

Introduction

Sepsis is a systemic inflammatory response caused by infection, trauma, and other factors, resulting in multi-organ dysfunction. In severe cases, it can progress to life-threatening septic shock and organ failure [1, 2]. Sepsis mainly induces liver injury, lung injury, cardiac dysfunction, gastrointestinal injury, and kidney injury. Among them, the liver is the most important

detoxification and metabolic organ of the human body, and it is very susceptible to infection and damage. Previous studies have shown that the reasons for the pathogenesis of sepsis-associated liver injury include oxidative stress (OS), hepatic microcirculation disorders, energy metabolism disorders, inflammatory response, etc. [3-5]. When sepsis occurs, the main mechanism of liver injury is OS due to increased oxygen free radicals and peroxidation of lipids

Address correspondence to: Shangping Fang, Anesthesia Laboratory and Training Center, School of Anesthesiology, Wannan Medical College, No. 22, Wenchang West Road, Lugang Street, Yijiang District, Wuhu 241002, Anhui, China. Tel: 19855362767; E-mail: 20180041@wnmc.edu.cn.

in the body [6]. Based on a review of recent literature, this paper summarizes the mechanism of OS-induced liver injury in sepsis, and the signaling pathways of reactive oxygen species (ROS) in sepsis-associated liver injury, as well as the effects of ROS on hepatocyte injury and antioxidant therapy.

Review progress

Oxidative stress

Oxidative damage

OS is a state of imbalance between oxidant and antioxidant in the body. When ROS and reactive nitrogen species such as nitric oxide (NO) increase in large quantities, the absence/shortage of antioxidants disrupts the balance in the body [7, 8]. ROS refers to oxygen free radicals and non-free radicals produced by the body, mainly containing the substrates of the respiratory chain of the inner mitochondrial membrane, such as $O_2^{\cdot -}$, $HO^{\cdot}$ and H_2O_2 , with oxidizing ability and toxic effects on most cells [9]. In a physiological state, as a by-product of the oxidation system, the production and clearance of ROS are in dynamic equilibrium in the body. Additionally, the level of ROS rises and exceeds the body's clearance ability, which can increase lipid peroxidation in related tissues, leading to abnormal expression of related proteins, DNA damage, and gene mutation [10]. Malondialdehyde (MDA), a lipid free radical, is the principal degradation product of lipid peroxides, which can damage mitochondrial and cell membranes, thereby inhibiting the function of membrane proteins, impairing the structural integrity of mitochondria, and leading to cell damage [10]. At the same time, NO synthase is greatly augmented, and NO can restrain the binding of mitochondrial cytochrome C to O_2 , leading to mitochondrial dysfunction. Thus, damage to cellular mitochondria causes spillover of cytochrome C from mitochondria into the cytoplasm, contributing to apoptosis [11, 12]. Through a series of reactions, it eventually contributes to autophagy, necrosis, and apoptosis, etc.

Anti-oxidation system

The antioxidant system is composed of endogenous molecules, including both enzymes and non-enzymes. The main enzymes include superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and catalase (CAT). To put it more precisely, SOD is an important antioxidant enzyme mainly involved in scavenging oxygen radicals. When ROS increases, SOD reacts with

$O_2^{\cdot -}$ to produce H_2O_2 , which is then converted into water by CAT and GSH-Px to remove ROS and protect cells from damage [13]. In contrast, when the contents of CAT, GSH-Px and SOD increase, the ROS and the MDA in the body reduce. The non-enzymes include vitamin C, vitamin E, uric acid, metal-binding protein, polyamines, bilirubin, carotenoids, etc. [14].

OS-related signaling pathways

Nuclear erythrocytes 2-related factor (Nrf2) signaling pathway

Nrf2 is a key factor in cellular oxidative stress and is negatively regulated by cytoplasmic Kelch-like ECH-associated protein 1 (Keap1) [15]. Under normal physiological conditions, Keap1 is a substrate linker protein of the Cul3 (Cul3)-dependent E3 ubiquitin ligase complex. Keap1 contains three functional domains, a Broad-complex, Tramtrack, and Bric-à-brac (BTB) domain, an intervening region domain, and a Kelch or double-glycine repeat domain. Wherein, BTB domain binds to Cul3 to assemble into a functional E3 ubiquitin ligase complex (Keap1-Cul3-E3). Keap1-Cul3-E3 ubiquitin ligase targets the Neh2 domain located at the N-terminus of Nrf2 and promotes Nrf2 ubiquitination [16]. After that, ubiquitinated Nrf2 is delivered to the 26S proteasome for degradation.

When the body is in an OS state, ROS modifies specific cysteine residues in Keap1, causing conformational changes in Keap1-Cul3-E3 ubiquitin ligase and interfering with Nrf2 ubiquitination [17]. Alternatively, it facilitates the breakdown of Nrf2 with Keap1 by activating protein kinase C. Nrf2 is then transferred into the nucleus by oxidative stressors such as electrophiles and binds to the antioxidant response element (ARE) by heterodimerization with the small Maf transcription factors [18]. Subsequently, the Nrf2-ARE signaling pathway initiates and promotes the expression of antioxidant enzyme genes, such as CAT, SOD, GSH-Px, glutathione S-transferase, NADH quinone oxidoreductase 1, and heme oxygenase 1, which in turn provides antioxidant defense [19, 20]. Moreover, ROS activates glycogen synthase kinase 3, nucleating phosphorylated Nrf2 and Keap1 and affecting the degradation of Nrf2 (Figure 1) [21]. In general, ROS can promote Nrf2 into the nucleus through various ways, and then activate its downstream pathway.

Nuclear factor kappa-light-chain-enhancer of activated B cells (NF-κB) signaling pathway

NF-κB is a transcription factor protein that con-




Figure 1. Effects of ROS on different signaling pathways. ROS, reactive oxygen species; Keap1, Kelch-like ECH-associated protein 1; Cul3, Cullin3; Nrf2, Nuclear erythrocytes 2-related factor; ARE, antioxidant response element; Ubc, E2-ubiquitin conjugating enzyme; NEDD8, neural precursor cell expressed developmentally downregulated 8; NLRP3, Nucleotide-binding oligomerization domain-like receptor protein 3; NIK, Nuclear factor kappa-light-chain-enhancer of activated B cells inducing kinase; IKK, IκB kinase; MEKK1, mitogen-activated protein kinase kinase kinase 1; AMPK, adenosine 5'-monophosphate-activated protein kinase; PI3K, Phosphatidylinositol 3-kinase; PIP3, phosphatidylinositol 3, 4, 5-triphosphate; AKT, activate protein kinase B; TSC, tuberous sclerosis complex (consisting of TSC1, TSC2, TBC1D7); VEGF, vascular endothelial growth factor; PTEN, phosphatase and tensin homolog; RTK, receptor tyrosine kinase; shc, recombinant protein; GRB, growth factor receptor-bound protein; SOS, son of sevenless; PLCγ, specific phospholipase Cγ; STAT, Signal transducer and activator of transcription; SOCS, suppressor of cytokine signaling; SHP-2, Src homology 2 domain-containing protein tyrosine phosphatase; JAK, janus kinase; MAPK, Mitogen-activated protein kinase; MEF2, myocyte enhancer factor 2 (contain: MEF2A, MEF2C, MEF2D); ELK1, ETS transcription factor; ATF-2, activating transcription factor 2; ERK, extra-cellular signal-regulated kinase; ASK1, apoptosis signal-regulating kinase 1; PDGF, platelet-derived growth factor; JNK, c-Jun N-terminal kinase; PPAR, Peroxisome proliferator-activated receptor; HIF-1α, hypoxia-inducible factor-1 alpha; TXNIP, Thioredoxin-interacting protein; GSH, glutathione; PKC, protein kinase C; Raf, rapidly accelerated fibrosarcoma; OX, oxide. → : activation; —|: inhibition.

tains five different family proteins: Rel A (p65), Rel B, c- ϵ l, p105 (p50 precursor/NF- κ B1), and p100 (p52 precursor/ NF- κ B2), by forming homologous or heterologous dimers with the DNA-binding protein -Rel [22]. NF- κ B is regulated by proteins of the I κ Bs family, including I κ B α , I κ B β , I κ B γ , I κ B ϵ , and B cell leukemia-3. In a physiological state, NF- κ B is inactivated in the cytoplasm due to its binding with I κ B α [23]. NF- κ B activation occurs through two major signaling pathways: canonical and non-canonical NF- κ B signaling pathways. The canonical pathway refers to the activation of I κ B kinase (IKK) and subsequent phosphorylation of I κ B α . I κ B α in the complex dissociates from NF- κ B1, activating NF- κ B1 [24]. The non-canonical NF- κ B pathway refers to NF- κ B inducing kinase activation and subsequent phosphorylation of IKK α . IKK α phosphorylates p100 to degrade to p52 [25]. Following this, the combination of Rel B and p52 translocates to the nucleus to regulate

gene transcription [26].

During OS, ROS mainly affects the activation of NF- κ B by activating phosphorylation of I κ B α and adjusting the mitogen-activated protein kinase kinase kinase 1 (MEKK1) upstream of IKK [27-29]. ROS can also interfere with ubiquitination and degradation of I κ B by inactivating Ubc12, thereby impacting the activation of NF- κ B [30]. In the non-canonical pathway, ROS could activate NF- κ B2 by upregulating NF- κ B inducing kinase (Figure 1) [31].

Nucleotide-binding oligomerization domain-like receptor protein 3 (NLRP3) signaling pathway

NLRP3 can form NLRP3 inflammasome bodies with precursors of apoptosis-associated spot-like protein in the caspase recruitment domain and cysteinyl aspartate specific proteinase 1 (Caspase-1). NLRP3 inflammasome is a

multi-protein macromolecular complex that processes Pro Caspase-1 into bioactive Caspase-1, cuts Gasdermin protein to cause cell perforation and mediates interleukin (IL) 1 β - and IL-18-related responses.

With the invasion of pathogen-associated molecular patterns or damage associated molecular patterns, mitochondria are stimulated to produce ROS. ROS activates NLRP3 inflammasomes or reacts with thioredoxin interaction, which then releases thioredoxin and activates NLRP3 inflammasomes (**Figure 1**) [32].

Mitogen-activated protein kinase (MAPK) signaling pathway

The cascade of MAPK consists of four main components: extracellular signal-associated kinase (ERK1/2), c-Jun N-terminal kinase, p38 kinase (p38), and large mitogen-activated protein (MAP) kinase 1 (BMK1/ERK5) [33]. MAP kinase kinase kinase (MAPKKK/MEKK/MKK) is activated, which then phosphorylates and activates MAP kinase kinase (MAPKK/MEK/MKK). Phosphorylation of MAPKK activates MAP kinase (MAPK), which, in turn, phosphorylates various substrate proteins [34].

ROS can activate receptors of epidermal growth factor and platelet-derived growth factor, activating Ras on cell membranes. Ras then recruits rapidly accelerated fibrosarcoma (Raf) (MAPKKK) in cytoplasm into the cell membrane for activation [35]. Activated Raf phosphorylates MEK1/2(MAPKK), which then results in ERK activation [36]. ROS acts on thioredoxin and glutaraldehyde, dissociating them from MAPKKKs such as apoptosis signal-regulating kinase 1 and MEKK1-4 [37]. This activation of MAPKKKs leads to the activation of MAPKK, subsequently activating c-Jun N-terminal kinase, p38, and BMK1 (**Figure 1**).

Phosphatidylinositol 3-kinase (PI3K) signaling pathway

PI3K catalyzes the synthesis of phosphatidylinositol 3, 4, 5-triphosphate to recruit and activate protein kinase B (AKT). Furthermore, it promotes the activation and transcription of Sestrins, endothelial nitric oxide synthase, forkhead box protein O, mechanistic target of rapamycin 1, and p53.

ROS can not only activate PI3K and its downstream signaling pathways, but also inactivate phosphatase and tensin homolog (PTEN) and inhibit AKT activation by negatively regulating phosphatidylinositol 3, 4, 5-triphosphate. Its

downstream pathway, Sestrins, comprises stress response proteins crucial for cellular adaptation to OS. Sestrins play a significant role in enhancing the body's tolerance to OS, preventing the accumulation of ROS and MDA through its inherent antioxidant activity and the regeneration of thiol peroxidase (peroxiredoxin) [38, 39]. Sestrin2 also inhibits sepsis-associated liver injury by activating the transcription factors p53, Nrf2, and hypoxia-inducible factor-1 α [40]. The other downstream pathway, mTOR, is a class of serine/threonine kinases that affect the OS of the body by regulating hypoxia-inducible factor-1 α , peroxisome proliferators-activated receptors- γ and other pathways (**Figure 1**).

Signal transducer and activator of transcription (STAT) signaling pathway

STAT, including STAT1-4, STAT5 (STAT5A and STAT5B), and STAT6, is also an important protein for OS. When IL-6 and other cytokines bind to the relevant receptors, they can change the molecular conformation of the receptor [41]. The coupled Janus kinase (JAK) kinases approach each other and phosphorylate tyrosine to activate the JAK kinases. The activated JAK kinases allow the corresponding STATs to form docking site with the receptor complex and bind to the receptor through its SH2 domain [42]. In response to JAK kinase, STATs are activated by phosphorylation and then form homo/heterodimers for incorporation into the nucleus. Dimeric STATs can bind to the promoters of target genes, and then the transcription of related genes can be activated [43].

During OS, ROS can inactivate tyrosine phosphatases and STAT phosphorylation. STAT3 also up-regulates glutathione to scavenge ROS (**Figure 1**) [44, 45].

Molecular mechanism of OS in sepsis-associated liver injury and antioxidant therapy

Hepatocytes

The roles of ROS production and accumulation in hepatocyte membrane and OS in sepsis-associated liver injury should not be underestimated. The main site of OS damage is the hepatocyte membrane. ROS can degrade unsaturated fatty acids on the hepatocyte membrane into lipid peroxides, which can aggravate the damage to mitochondria [46]. After mitochondria are damaged by ROS, they release cytochrome C into the cytoplasm and cause hepatocyte apoptosis. When sepsis occurs, the content of inducible nitric oxide synthase in the liver is significantly increased. Inducible nitric

Table 1. Signaling pathways and therapeutics of different cellular antioxidant therapies in liver

Types	Signaling pathway	Therapeutic agents
Hepatocyte	mTOR / p70S6K	Albiflorin [53], Anemoside B4 [54]
	Nrf2 / NF-κB	Wogonin [52]
	SIRT1 / STAT3	Melatonin [55]
	SIRT1 / AMPK	Ginsenoside R3 [51]
	Nrf2 / NLRP3	Malvidin [50]
	RAGE / p38	Idebenone [49]
Hepatic stellate cells	VEGF	Dihydroartemisinin [62]
	Nrf-2	Schisandrin B [60]
	ZEB-2 / PTEN	Orientin [59]
	CYP2E/Nrf2 / ROS and PI3K / AKT	Sophocarpine [63]
	ROS / NF-κB	Chlorogenic acid [64]
Hepatic sinusoidal endothelial cells	TLR4 / NFκB	Liquiritigenin and liquiritin [68]
	ROS / ERK	Curcumol [67]
Kupffer cells	CUL3	Pyrroloquinoline quinone [71]
	AMPK	Isoquercetin [72]
	TLR4 / NF-κB / NLRP3	Ginsenoside Rb1 [73]

Note: Nrf2, Nuclear erythrocytes 2-related factor; p70S6K, ribosomal protein S6 kinase; NF-κB, Nuclear factor kappa-light-chain-enhancer of activated B cells; SIRT1, silent information regulator 1; STAT, signal transducer and activator of transcription; AMPK, adenosine 5'-monophosphate-activated protein kinase; NLRP3, Nucleotide-binding oligomerization domain-like receptor protein 3; RAGE, receptor for advanced glycation end products; VEGF, vascular endothelial growth factor; ZEB-2, zinc finger E-box binding homeobox2; PTEN, phosphatase and tensin homolog; CYP2E1, cytochrome P450 family 2 subfamily E member 1; ROS, reactive oxygen species; ERK, extra-cellular signal-regulated kinase; PI3K, Phosphatidylinositol 3-kinase; AKT, activate protein kinase B; TLR4, toll-like receptor 4; Cul3, Cullin3.

oxide synthase can induce a large amount of NO, which in turn can react with mitochondrial cytochrome oxidase to inhibit the activity of cytochrome oxidase, and then bring about structural and functional damage on mitochondrial proteins. NO can also react with O²⁻ to generate strong oxidative nitrite peroxide ions, which can oxidize with various molecules, giving rise to apoptosis [47].

Celep et al. found that silymarin could alleviate sepsis-associated liver injury by reducing OS [48]. Gou et al. revealed that Idebenone could prevent sepsis-induced OS, apoptosis, and inflammation by inhibiting RAGE/p38 signaling [49]. Fan et al. discovered that Malvidin prevented the lipopolysaccharide (LPS)-induced decrease in antioxidant enzyme activity by upregulating the Nrf2 pathway, and at the same time inhibited the NLRP3 inflammasome, apoptosis, and autophagy, thereby alleviating sepsis-associated liver injury [50]. Wu et al. reported that ginsenoside Rg3 decreased ROS and stimulated the SIRT1/AMPK pathway, thereby enhancing autophagy to ameliorate sepsis-associated liver injury and mitochondrial dysfunction [51]. Dai et al. demonstrated that Wogonin improved both OS and inflammatory response by activating Nrf2, thus acting as a protective agent against sepsis-associated

liver injury [52]. Liu et al. and Pei et al. found that both albiflorin and anemoside B4 inhibited LPS-induced apoptosis and OS of primary hepatocytes through the mTOR/p70S6K signaling pathway, and alleviated liver damage induced by sepsis [53, 54]. Chen et al. and Kleber et al. concluded that melatonin played a protective role in sepsis-induced liver injury by reducing ROS and activating SIRT1/STAT3 pathway [55, 56] (**Table 1**).

Hepatic stellate cells (HSCs)

HSCs are endothelial pericytes located in the hepatic sinusoid space. During liver injury, HSCs are activated into myofibroblasts and secrete extracellular matrix to participate in injury repair. At the same time, the activation of HSCs is also the central link in the development and progression of liver fibrosis [57]. HSCs also bring forth a large number of cytokines, chemokines, and growth mediators and express cell adhesion molecules in response to stimuli such as endotoxin (LPS). Hepatocyte injury results in the release of damage associated molecular patterns, which stimulates HSCs and Kupffer cells (KCs). Further, stimulation of HSCs and KCs recruit immune cells and secrete proinflammatory mediators. Interaction between HSCs and KCs produces a two-way environmen-

tal interaction, leading to the release of pro-inflammatory and anti-inflammatory cytokines. This continued release contributes to ongoing liver damage [58].

It has been found that orientin inhibited HSC activation and reduced LPS and CCl₄-induced liver injury by down-regulating ZEB-2/PTEN [59]. Yao et al. found that schisandrin B attenuated OS in LPS-induced HSC activation through the Nrf2 signaling pathway [60]. Rani et al. revealed that HSCs-sufficient mice treated with acetaminophen exhibited stronger OS compared to HSCs-depleted mice, suggesting that HSCs plays a major role in OS in hepatocytes [61]. Huan et al. found that dihydroartemisinin inhibited HSCs activation by regulating miR-29b-3p to affect vascular endothelial growth factor pathway [62]. Jiang et al. reported that sophocarpine alleviated LPS-induced liver injury, OS and inflammation by inhibiting CYP2E/Nrf2/ROS and PI3K/AKT pathways [63]. Shi et al. demonstrated that chlorogenic acid inhibited LPS-induced proinflammatory responses to HSCs through inhibition of the ROS/NF-κB signaling pathway (**Table 1**) [64].

Hepatic sinusoidal endothelial cells

Liver sinusoidal endothelial cells (LSECs) are highly sensitive to OS. ROS and reactive nitrogen species can reduce the activity of endothelial nitric oxide synthase messenger RNA, leading to dysfunction of LSECs. This dysfunction promotes vasoconstriction, increases intercellular block, raises vascular permeability, and accelerates thrombosis, and further aggravates liver injury [65].

Research indicated that artesunate enhanced the defense of LSECs and prevented the aggregation of inflammatory immune cells, mitigating the effects of sepsis-associated liver injury, inflammation, and dysfunction [66]. Gao et al. demonstrated that curcumin inhibited the angiogenesis of recombinant Kruppel like factor 5-mediated LSECs by inhibiting ROS/ERK signaling [67]. Chen et al. found that glycyrrhizin attenuated cyclophosphamide-induced LSEC injury and inflammatory injury in mice by reducing liver peroxidase activity (**Table 1**) [68]. Pioglitazone can suppress the activation of NF-κB, activator protein-1, MAPK and other inflammatory signaling pathways, and inhibit the release of ROS by KCs, the generation of cytokines and pro-inflammatory mediators, and the expression of adhesion molecule genes in LSECs. Tai et al. discovered that celecoxib reduced ROS through both cyclooxygenase-2 dependent and independent signaling pathways. This reduction

in ROS is associated with a significant decrease in NO clearance, subsequently reducing LSEC capillaries and correcting endothelial dysfunction [69].

KCs

KCs are the most important resident macrophages in the liver with abilities of self-renewal, clearance of endotoxin by phagocytosis and release of inflammatory factors to maintain immune homeostasis in the liver. At the same time, during sepsis, KCs can facilitate the development of OS by releasing ROS. KCs further interact with hepatocytes to produce and release chemokines to recruit monocytes in the circulation and subsequently differentiate into monocyte-derived macrophages [70].

Wu et al. found that pyrroloquinoline quinone could inhibit Cul3 expression to alleviate OS and inflammation caused by KCs in sepsis-associated liver injury [71]. Qin et al. revealed that Isoquercetin alleviated inflammation and OS by activating AMPK pathway and inhibiting transforming growth factor β signaling [72]. Liu et al. found that ginsenoside Rb1 attenuated OS and inflammation through TLR4/NF-κB signaling pathway and NLRP3 inflammasome activation [73]. Zhou et al. demonstrated that ultralow doses of dexamethorphan reduced superoxide radicals and alleviated OS in KCs (**Table 1**) [74]. Zhai et al. reported that NADPH oxidase 4 activated the NLRP3 inflammasome through the generation of ROS. This activation promotes the inflammatory response of KC and the release of inflammatory factors, and inhibition of NADPH oxidase 4 ameliorated acute liver injury in mice [75]. Zhang et al. observed that stimulator of interferon genes signaling in KCs enhanced ROS production in response to LPS stimulation, which upgraded mitochondrial DNA release and hepatocyte death, contributing to LPS-induced liver injury [76].

Conclusion

In summary, OS plays an important role in the development and progression of sepsis-associated liver injury. Nrf2, NF-κB, PI3K, NLRP3, MAPK, STAT and other signaling pathways can regulate OS. This review provides the theoretical basis for the antioxidant therapy of sepsis-associated liver injury. Furthermore, liver circulatory and metabolic disorders, excessive inflammatory response, mitochondrial dysfunction, neutrophil and neutrophil extracellular trapping, and autophagy are all mechanisms leading to sepsis associated liver injury. At present, there are many studies on hepatic paren-

chymal cells and KCs in sepsis-associated liver injury. Most of the studies on hepatic stellate cells and hepatic sinusoidal endothelial cells are related to the treatment of liver fibrosis and cirrhosis, but few on sepsis-associated liver injury. This paper summarizes the OS status of different hepatocytes and the treatment of parenchymal and non-parenchymal cells, which can provide ideas for basic clinical research on sepsis related liver injury, so as to reduce the mortality of sepsis related liver injury.

References

- [1] Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *Jama* 2016;315:801-810.
- [2] Zhu B. Recent advances in the research of viral sepsis. *Zhonghua Er Ke Za Zhi* 2021;59:250-253.
- [3] Zhao GM, Hu ZS. Research progress on the pathogenesis and treatment of sepsis-induced liver injury. *Jiefangjun Yixue Zazhi* 2019;44:515-520.
- [4] Sakr Y, Dubois MJ, De Backer D, et al. Persistent microcirculatory alterations are associated with organ failure and death in patients with septic shock. *Crit Care Med* 2004;32:1825-1831.
- [5] Mainali R, Zabalawi M, Long D, et al. Dichloroacetate reverses sepsis-induced hepatic metabolic dysfunction. *Elife* 2021;10:e64611.
- [6] Sakaguchi S, Furusawa S. Oxidative stress and septic shock: metabolic aspects of oxygen-derived free radicals generated in the liver during endotoxemia. *FEMS Immunol Med Microbiol* 2006;47:167-177.
- [7] Xu F, Xu J, Xiong X, et al. Salidroside inhibits MAPK, NF- κ B, and STAT3 pathways in psoriasis-associated oxidative stress via SIRT1 activation. *Redox Rep* 2019;24:70-74.
- [8] Diesen DL, Kuo PC. Nitric oxide and redox regulation in the liver: part II. Redox biology in pathologic hepatocytes and implications for intervention. *J Surg Res* 2011;167:96-112.
- [9] Pizzino G, Irrera N, Cucinotta M, et al. Oxidative Stress: Harms and Benefits for Human Health. *Oxid Med Cell Longev* 2017;2017:8416763.
- [10] Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev* 2014;2014:360438.
- [11] Lu JQ, Liu JY, Li A. Roles of neutrophil reactive oxygen species (ROS) generation in organ function impairment in sepsis. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)* 2022;23:437-451.
- [12] Borutaite V, Morkuniene R, Arandarcikaite O, et al. Nitric oxide protects the heart from ischemia-induced apoptosis and mitochondrial damage via protein kinase G mediated blockage of permeability transition and cytochrome c release. *J Biomed Sci* 2009;16:1-12.
- [13] Moris D, Spartalis M, Tzatzaki E, et al. The role of reactive oxygen species in myocardial redox signaling and regulation. *Ann Transl Med* 2017;5:324.
- [14] Zhang Z, Rong L, Li YP. Flaviviridae viruses and oxidative stress: implications for viral pathogenesis. *Oxid Med Cell Longev* 2019;2019:1-17.
- [15] Itoh K, Wakabayashi N, Katoh Y, et al. Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 1999;13:76-86.
- [16] Zhang DD, Lo S, Cross J, et al. Keap1 Is a Redox-Regulated Substrate Adaptor Protein for a Cul3-Dependent Ubiquitin Ligase Complex. *Mol Cell Biol* 2004;24:10941-10953.
- [17] Sajadimajd S, Khazaei M. Oxidative stress and cancer: the role of Nrf2. *Curr Cancer Drug Targets* 2018;18:538-557.
- [18] Otsuki A, Yamamoto M. Cis-element architecture of Nrf2-sMaf heterodimer binding sites and its relation to diseases. *Arch Pharm Res* 2020;43:275-285.
- [19] Mann GE, Niehueser-Saran J, Watson A, et al. Nrf2/ARE regulated antioxidant gene expression in endothelial and smooth muscle cells in oxidative stress: implications for atherosclerosis and preeclampsia. *Acta Physiol Sin* 2007;59:117-127.
- [20] Lee J-M, Johnson JA. An important role of Nrf2-ARE pathway in the cellular defense mechanism. *J Biochem Mol Biol* 2004;37:139-143.
- [21] Zhang J, Wang X, Vikash V, et al. ROS and ROS-mediated cellular signaling. *Oxid Med Cell Longev* 2016;2016:4350965.
- [22] Morgan MJ, Liu ZG. Crosstalk of reactive oxygen species and NF-kappaB signaling. *Cell Res* 2011;21:103-115.
- [23] Ito CY, Kazantsev AG, Baldwin AS. Three NF-kappa B sites in the I kappa B-alpha promoter are required for induction of gene expression by TNF alpha. *Nucleic Acids Res* 1994;22:3787-3792.
- [24] Beinke S, Ley SC. Functions of NF- κ B1 and NF- κ B2 in immune cell biology. *Biochem J* 2004;382:393-409.
- [25] Xiao G, Harhaj EW, Sun SC. NF- κ B-inducing kinase regulates the processing of NF- κ B2 p100. *Mol Cell* 2001;7:401-409.
- [26] Sun SC. The non-canonical NF- κ B pathway in

- immunity and inflammation. *Nat Rev Immunol* 2017;17:545-558.
- [27] Tilstra JS, Gaddy DF, Zhao J, et al. Pharmacologic IKK/NF-kappa B inhibition causes antigen presenting cells to undergo TNF alpha dependent ROS-mediated programmed cell death. *Sci Rep* 2014;4:3631.
- [28] Cross J, Templeton D. Oxidative stress inhibits MEKK1 by site-specific glutathionylation in the ATP-binding domain. *Biochem J* 2004;381:675-683.
- [29] Janssen-Heininger YM, Macara I, Mossman BT. Cooperativity between oxidants and tumor necrosis factor in the activation of nuclear factor (NF)-kappa b: requirement of ras/mitogen-activated protein kinases in the activation of NF-kappa B by oxidants. *Am J Respir Cell Mol Biol* 1999;20:942-952.
- [30] Zhang ZX, Heng YQ, Cheng W, et al. Reactive oxygen species (ROS)-response nanomedicine through knocking down a novel therapeutic target NEDD8-conjugating enzyme UBC12 (UBE2M) in the treatment of liver cancer. *Mater Des* 2021;204:14.
- [31] Zhong X, Zhang Z, Shen H, et al. Hepatic NF-kappa B-Inducing Kinase and Inhibitor of NF-kappa B Kinase Subunit alpha Promote Liver Oxidative Stress, Ferroptosis, and Liver Injury. *Hepatol Commun* 2021;5:1704-1720.
- [32] Zhou RB, Tardivel A, Thorens B, et al. Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat Immunol* 2010;11:136-140.
- [33] Cui X, Qian DW, Jiang S, et al. Scutellariae Radix and Coptidis Rhizoma Improve Glucose and Lipid Metabolism in T2DM Rats via Regulation of the Metabolic Profiling and MAPK/PI3K/Akt Signaling Pathway. *Int J Mol Sci*. 2018;19(11):3634.
- [34] Morigasaki S, Ikner A, Tatebe H, et al. Response regulator-mediated MAPKKK heteromer promotes stress signaling to the Spc1 MAPK in fission yeast. *Mol Biol Cell* 2013;24:1083-1092.
- [35] Franklin RA, Tordai A, Patel H, et al. Ligation of the T cell receptor complex results in activation of the Ras/Raf-1/MEK/MAPK cascade in human T lymphocytes. *J Clin Invest* 1994;93:2134-2140.
- [36] Williams NG, Paradis H, Agarwal S, et al. Raf-1 and p21v-ras cooperate in the activation of mitogen-activated protein kinase. *Proc Natl Acad Sci USA* 1993;90:5772-5776.
- [37] Tobiume K, Matsuzawa A, Takahashi T, et al. ASK1 is required for sustained activations of JNK/p38 MAP kinases and apoptosis. *EMBO Rep* 2001;2:222-228.
- [38] Haidurov A, Budanov AV. Sestrin family-the stem controlling healthy ageing. *Mech Ageing Dev* 2020;192:111379.
- [39] Kim H, An S, Ro SH, et al. Janus-faced Sestrin2 controls ROS and mTOR signalling through two separate functional domains. *Nat Commun* 2015;6:10025.
- [40] Park SJ, Cho SS, Kim KM, et al. Protective effect of sestrin2 against iron overload and ferroptosis-induced liver injury. *Toxicol Appl Pharmacol* 2019;379:114665.
- [41] Luticken C, Wegenka UM, Yuan J, et al. Association of transcription factor APRF and protein kinase Jak1 with the interleukin-6 signal transducer gp130. *Science* 1994;263:89-92.
- [42] Schaper F, Gendo C, Eck M, et al. Activation of the protein tyrosine phosphatase SHP2 via the interleukin-6 signal transducing receptor protein gp130 requires tyrosine kinase Jak1 and limits acute-phase protein expression. *Biochem J* 1998;335:557-565.
- [43] Williamson L, Ayalon I, Shen H, et al. Hepatic STAT3 inhibition amplifies the inflammatory response in obese mice during sepsis. *Am J Physiol Endocrinol Metab* 2019;316:E286-E292.
- [44] Linher-Melville K, Singh G. The complex roles of STAT3 and STAT5 in maintaining redox balance: Lessons from STAT-mediated xCT expression in cancer cells. *Mol Cell Endocrinol* 2017;451:40-52.
- [45] Zhang W, Gong M, Zhang W, et al. Thiostrepton induces ferroptosis in pancreatic cancer cells through STAT3/GPX4 signalling. *Cell Death Dis* 2022;13:630.
- [46] Schenk B, Fulda S. Reactive oxygen species regulate Smac mimetic/TNFalpha-induced necroptotic signaling and cell death. *Oncogene* 2015;34:5796-5806.
- [47] Campolo N, Bartesaghi S, Radi R. Metal-catalyzed protein tyrosine nitration in biological systems. *Redox Rep* 2014;19:221-231.
- [48] Celep NA, Gedikli S. Protective Effect of Silymarin on Liver in Experimental in the Sepsis Model of Rats. *Acta Histochem Cytochem* 2023;56:9-19.
- [49] Gou T, Jin X, Xia J. Idebenone reduces sepsis-induced oxidative stress and apoptosis in hepatocytes via RAGE/p38 signaling. *Ann Transl Med* 2022;10:1363.
- [50] Fan H, Cui J, Liu F, et al. Malvidin protects against lipopolysaccharide-induced acute liver injury in mice via regulating Nrf2 and NLRP3 pathways and suppressing apoptosis and autophagy. *Eur J Pharmacol* 2022;933:175252.
- [51] Wu P, Yu X, Peng Y, et al. Ginsenoside Rg3 alleviates septic liver injury by regulating the lncRNA TUG1/miR-200c-3p/SIRT1 axis. *J Inflamm (Lond)* 2021;18:31.

- [52] Dai JM, Guo WN, Tan YZ, et al. Wogonin alleviates liver injury in sepsis through Nrf2-mediated NF-kappa B signalling suppression. *J Cell Mol Med* 2021;25:5782-5798.
- [53] Liu Y, Feng L, Yao L. Albiflorin alleviates sepsis-induced acute liver injury through mTOR/p70S6K pathway. *Curr Mol Med* 2023.
- [54] Pei L, He L. Hepatoprotective effect of anemoside B4 against sepsis-induced acute liver injury through modulating the mTOR/p70S6K-mediated autophagy. *Chem Biol Interact* 2021;345:109534.
- [55] Chen J, Xia H, Zhang L, et al. Protective effects of melatonin on sepsis-induced liver injury and dysregulation of gluconeogenesis in rats through activating SIRT1/STAT3 pathway. *Biomed Pharmacother* 2019;117:109150.
- [56] Kleber A, Kubulus D, Rössler D, et al. Melatonin modifies cellular stress in the liver of septic mice by reducing reactive oxygen species and increasing the unfolded protein response. *Exp Mol Pathol* 2014;97:565-571.
- [57] Tsuchida T. Mechanisms of hepatic stellate cell activation as a therapeutic target for the treatment of non-alcoholic steatohepatitis. *Nihon Yakurigaku Zasshi* 2019;154:203-209.
- [58] Mooli RGR, Mukhi D, Ramakrishnan SK. Oxidative stress and redox signaling in the pathophysiology of liver diseases. *Compr Physiol* 2022;12:3167-3192.
- [59] Khalil HE, Ibrahim H-IM, El-Fass KA, et al. Orientin Alleviates Liver Inflammation via Downregulation of ZEB-2/PTEN Markers-Hepatic Stellate Cells Approach. *Appl Sci -Basel* 2022;12:2725.
- [60] Yao Z, Han J, Lou S, et al. Schisandrin B attenuates lipopolysaccharide-induced activation of hepatic stellate cells through Nrf-2-activating anti-oxidative activity. *Int J Clin Exp Pathol* 2018;11:4917-4925.
- [61] Rani R, Sharma A, Wang J, et al. Endotoxin-stimulated hepatic stellate cells augment acetaminophen-induced hepatocyte injury. *Am J Pathol* 2022;192:518-535.
- [62] Huan S, Sun S, Song S, et al. Dihydroartemisinin inhibits the activation and proliferation of hepatic stellate cells by regulating miR-29b-3p. *Int J Mol Med* 2023;51:1-16.
- [63] Jiang Z, Meng Y, Bo L, et al. Sophocarpine Attenuates LPS-Induced Liver Injury and Improves Survival of Mice through Suppressing Oxidative Stress, Inflammation, and Apoptosis. *Mediators Inflamm* 2018;2018:1-12.
- [64] Shi H, Dong L, Dang X, et al. Effect of chlorogenic acid on LPS-induced proinflammatory signaling in hepatic stellate cells. *Inflamm Res* 2013;62:581-587.
- [65] Lafoz E, Ruat M, Anton A, et al. The Endothelium as a Driver of Liver Fibrosis and Regeneration. *Cells* 2020;9:929.
- [66] He XL, Chen JY, Feng YL, et al. Single-cell RNA sequencing deciphers the mechanism of sepsis-induced liver injury and the therapeutic effects of artesunate. *Acta Pharmacol Sin* 2023;44:1801-1814.
- [67] Gao L, Yang X, Li Y, et al. Curcumin inhibits KLF5-dependent angiogenesis by blocking the ROS/ERK signaling in liver sinusoidal endothelial cells. *Life Sci* 2021;264:118696.
- [68] Chen M, Zhang C, Zhang J, et al. The involvement of DAMPs-mediated inflammation in cyclophosphamide-induced liver injury and the protection of liquiritigenin and liquiritin. *Eur J Pharmacol* 2019;856:172421.
- [69] Tai Y, Zhao C, Zhang L, et al. Celecoxib reduces hepatic vascular resistance in portal hypertension by amelioration of endothelial oxidative stress. *J Cell Mol Med* 2021;25:10389-10402.
- [70] Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med* 1989;320:365-376.
- [71] Wu Y, Zhao M, Lin Z. Pyrroloquinoline quinone (PQQ) alleviated sepsis-induced acute liver injury, inflammation, oxidative stress and cell apoptosis by downregulating CUL3 expression. *Bioengineered* 2021;12:2459-2468.
- [72] Qin G, Ma J, Huang Q, et al. Isoquercetin Improves Hepatic Lipid Accumulation by Activating AMPK Pathway and Suppressing TGF-beta Signaling on an HFD-Induced Nonalcoholic Fatty Liver Disease Rat Model. *Int J Mol Sci* 2018;19:4126.
- [73] Liu Y, Liu N, Liu Y, et al. Ginsenoside Rb1 Reduces D-GalN/LPS-induced Acute Liver Injury by Regulating TLR4/NF-kappa B Signaling and NLRP3 Inflammasome. *J Clin Transl Hepatol* 2022;10:474-485.
- [74] Zhou R, Chen SH, Li G, et al. Ultralow doses of dextromethorphan protect mice from endotoxin-induced sepsis-like hepatotoxicity. *Chem Biol Interact* 2019;303:50-56.
- [75] Zhai L, Pei H, Yang Y, et al. NOX4 promotes Kupffer cell inflammatory response via ROS-NLRP3 to aggravate liver inflammatory injury in acute liver injury. *Aging (Albany NY)* 2022;14:6905-6916..
- [76] Zhang Q, Wei J, Liu Z, et al. STING signaling sensing of DRP1-dependent mtDNA release in kupffer cells contributes to lipopolysaccharide-induced liver injury in mice. *Redox Biol* 2022;54:102367.